arXiv:1302.6963v2 [gr-qc] 1 Mar 2013

Numerical solutions to the cosmological 3-fluid problem

Mustapha Azreg-Aïnou
Başkent University, Department of Mathematics, Bağlıca Campus, Ankara, Turkey

Abstract

We show that, for the scalar field cosmology with exponential potential, the set of values of the coupling parameter for which the solutions undergo a transient period of acceleration is much larger than the set discussed in the literature. The gradual inclusion of ordinary and dark matters results in an everywhere, but near the origin, smoother and right shifted (along the time axis) acceleration curve. For the 3-fluid problem, the energy density need not exhibit a plateau during the acceleration period. Much excess in the dark matter and/or ordinary matter energy densities would lead the universe to undergo an eternal deceleration expansion.

PACS numbers: 98.80.-k, 95.35.+d, 95.36.+x, 04.25.D-

The problem. To account for the present period of acceleration of the universe many models have been suggested so far [1]- [15]. In this work we are concerned with the models corresponding to spatially flat Friedmann-Roberston-Walker (FRW) universes where the source is (1) a homogeneous scalar field coupled to exponential potential [representing dark energy (DE)] [8, 14] and references therein or (2) a homogeneous scalar field coupled to exponential potential along with the dark matter (DM), which is modeled by a dust fluid, and a perfect fluid representing ordinary matter (OM) [15] and references therein. This is the so-called 3-fluid problem.

The problem of a scalar field coupled to exponential potential has been solved in d-dimensional by decoupling the system of equations governing the dynamics [8]. If the potential is of the form $V = V_0 \mathrm{e}^{-\lambda\phi}$ where ϕ is the scalar field, the solutions have been classified according to the coupling constant λ [14]. The problem with a piecewise exponential potential in 4-dimensional has also been solved [14].

The 3-fluid problem that involves a scalar field (DE), a dust fluid (DM) and a perfect fluid has been treated analytically in [15]. The method developed in [15] and references therein that is based on symmetry assumptions has led, as was shown later in [16], to a flawed solution to the 3-fluid problem, in that the solution derived in [15] does not satisfy the field equations.

Thus, to our knowledge both problems of deriving analytic solutions or developing new methods to tackle to 3-fluid problem remain open. As we shall see below, even the linear case is not easy to handle. The purpose of this work is to provide numerical solutions to the 3-fluid problem with exponential potential and compare them to the solutions of the scalar field problem with similar potential.

To write explicitly the solutions for the normal (not phantom) scalar field problem we use a slightly different notation but the same conventions as in [14] with the exception of taking $V = V_0 \mathrm{e}^{-\lambda\phi}$ (instead of $V = V_0 \mathrm{e}^{+\lambda\phi}$). For the case of the scalar field alone we use the subscript s to denote any variable. For instance, we write $V_s = V_0 \mathrm{e}^{-\lambda\phi_s}$ and $H_s^2 = \rho_s$ for the scalar field problem and $V = V_0 \mathrm{e}^{-\lambda\phi}$ and $H^2 = \rho$ for the 3-fluid problem.

Consider the flat ($k = 0$) FRW cosmological model for the universe $\mathrm{d}s^2 = \mathrm{d}t^2 - a_s^2(t)\mathrm{d}\vec{x}^2$. The equations governing the motion of (a_s, ϕ_s) are the FRW equation $H_s^2 = \rho_s$ and the Klein-Gordon one, where

ρ_s is the energy density and $H_s = \dot{a}_s/a_s$ is the Hubble variable ($\dot{x} = \mathrm{d}x/\mathrm{d}t$)

$$\frac{\dot{a}_s{}^2}{a_s^2} = \frac{\dot{\phi}_s{}^2}{2} + V_0 \mathrm{e}^{-\lambda\phi_s} \tag{1}$$

$$\ddot{\phi}_s + 3\frac{\dot{a}_s}{a_s}\dot{\phi}_s - \lambda V_0 \mathrm{e}^{-\lambda\phi_s} = 0. \tag{2}$$

The solutions are called hyperbolic if $\lambda < 3\sqrt{2}$ or trigonometric if $\lambda > 3\sqrt{2}$ [14]. If we let $a_s = \mathrm{e}^{U_s/3}$, $\phi_s = \sqrt{2}W_s/3$, $\alpha = \sqrt{18-\lambda^2}/(3\sqrt{2})$, $\beta = \sqrt{\lambda^2-18}/(3\sqrt{2})$, and $\ell_\pm = 6/(6 \pm \sqrt{2}\lambda)$ the hyperbolic and trigonometric solutions are given by, respectively

$$U_s = 2\tau/\alpha + \ell_- \ln(1 + m\mathrm{e}^{-2\alpha\tau}) + \ell_+ \ln(1 - m\mathrm{e}^{-2\alpha\tau}) \tag{3}$$

$$W_s = \sqrt{2}\lambda\tau/(3\alpha) + \ell_- \ln(1 + m\mathrm{e}^{-2\alpha\tau}) - \ell_+ \ln(1 - m\mathrm{e}^{-2\alpha\tau}) \tag{4}$$

$$U_s = \ell_- \ln[\cos(\beta\tau)] + \ell_+ \ln[\sin(\beta\tau)] \tag{5}$$

$$W_s = \ell_- \ln[\cos(\beta\tau)] - \ell_+ \ln[\sin(\beta\tau)], \tag{6}$$

where we have omitted additive constants. Here m is a real constant and τ is the new time coordinate defined by

$$\dot{\tau} = \frac{3}{2}\sqrt{V_0}\mathrm{e}^{-\sqrt{2}\lambda W_s/6}. \tag{7}$$

If $m \neq 0$, then m may be set equal to ± 1 by shifting the values of τ. For all $\lambda < 3\sqrt{2}$, it is straightforward to show that the range of τ extends from a finite value (usually taken 0) to infinity, and for $\lambda < 3\sqrt{2}$, $0 < \tau < \pi/(2\beta)$.

The 3-fluid problem is also governed by the Klein-Gordon equation (2) (with the subscript s removed) and the FRW equation $H^2 = \rho$ with $H = \dot{a}/a$ and $\rho = \rho_\phi + \rho_{\mathrm{DM}} + \rho_{\mathrm{OM}}$ is the sum of the energy densities of the three fluids (scalar field, dark and ordinary matters). If the scalar field is normal, its energy density is given by

$$\rho_\phi = \frac{\dot{\phi}^2}{2} + V_0 \mathrm{e}^{-\lambda\phi}. \tag{8}$$

The DM is treated as a dust (without pressure) while the OM is a perfect fluid with an equation of state

$$p_{\mathrm{OM}} = (\gamma - 1)\rho_{\mathrm{OM}} \qquad (1 < \gamma \leq 2). \tag{9}$$

Using the three conservation equations (for each fluid $T^{\mu\nu}{}_{;\nu} = 0$ where $T^{\mu\nu}$ is the stress-energy tensor) and (9), one arrives at [15]

$$\rho_{\mathrm{OM}} = Ca^{-3\gamma}, \quad \rho_{\mathrm{DM}} = E/a^3, \tag{10}$$

where $C \geq 0$ and $E \geq 0$ are integration constants. The 3-fluid problem reduces to that of the scalar field if $C = E = 0$. The equations of motion take the form[1]

$$a\dot{a}^2 - \frac{1}{2}a^3\dot{\phi}^2 - V_0 a^3 \mathrm{e}^{-\lambda\phi} - Ca^{-3(\gamma-1)} = E \tag{11}$$

$$\ddot{\phi} + 3\frac{\dot{a}}{a}\dot{\phi} - \lambda V_0 \mathrm{e}^{-\lambda\phi} = 0. \tag{12}$$

Now we set $a = \mathrm{e}^{U/3}$ and $\phi = \sqrt{2}W/3$ reducing the system to

$$9E\mathrm{e}^{-U} = \dot{U}^2 - \dot{W}^2 - 9V_0\mathrm{e}^{-\sqrt{2}\lambda W/3} - 9C\mathrm{e}^{-\gamma U} \tag{13}$$

$$2\ddot{W} + 2\dot{U}\dot{W} - 3\sqrt{2}\lambda V_0 \mathrm{e}^{-\sqrt{2}\lambda W/3} = 0. \tag{14}$$

---

[1] The second term in Eq. (20) of Ref. [15] should read $-\varepsilon k a^3\dot{\phi}^2/2$.

As noticed earlier, if $C = E = 0$ the system (11) and (12) reduces to that of the scalar field problem, which is decoupled upon introducing the new variables (u_s, v_s) such that $U_s = v_s + u_s$ and $W_s = v_s - u_s$ and using the time τ defined in (7) [8]. It does not seem easy to decouple the system for $C \neq 0$ and $E \neq 0$, the case to which we are interested.

For the purpose of the numerical analysis we are aiming to perform, it is not convenient to introduce a new time coordinate τ defined by $\dot{\tau} = \frac{3}{2}\sqrt{V_0}\mathrm{e}^{-\sqrt{2}\lambda W/6}$. Rather, we will use the same time τ as defined in (7). This will allow us to compare solutions to both problems with the same initial conditions. The system (13) and (14) takes the form (with $x' = \mathrm{d}x/\mathrm{d}\tau$)

$$4E\mathrm{e}^{-U+\sqrt{2}\lambda W_s/3} = V_0U'^2 - V_0W'^2 - 4V_0\mathrm{e}^{-\sqrt{2}\lambda(W-W_s)/3} - 4C\mathrm{e}^{-\gamma U+\sqrt{2}\lambda W_s/3} \tag{15}$$

$$6W'' - \sqrt{2}\lambda W_s'W' + 6U'W' - 4\sqrt{2}\lambda\mathrm{e}^{-\sqrt{2}\lambda(W-W_s)/3}. \tag{16}$$

It is possible by a perturbation approach to decouple the system (15) and (16) if we assume that the independent constants C and E are small enough to allow for such an approach. In fact, writing $U = U_s + U_C C + U_E E + \cdots$ and $W = W_s + W_C C + W_E E + \cdots$ it is possible to determine the four time functions (U_C, U_E, W_C, W_E), however, their expressions are sizable and a numerical analysis is needed anyway. We would prefer applying a numerical approach to the full nonlinear system (15) and (16) where we can consider large values of C and E.

Numerical solutions. For the scalar field problem, it was noticed in Ref. [8] that the energy density exhibits a plateau during the transient period of acceleration (TPA). We verify that this statement remains valid for at least[2] $\sqrt{6} < \lambda < 3\sqrt{2}$ and $\lambda > 3\sqrt{2}$. However, the existence of a plateau for the energy density does not ensure the occurrence of a TPA: We have noticed the existence of solutions, as are the cases for $\lambda = \sqrt{2}$ and $\lambda = 2/\sqrt{3}$, with no TPA but (twice) the kinetic energy vanishes at the point where the energy density has a plateau and the potential energy is maximum. For instance, for $\lambda = \sqrt{2}$ a plot of the energy density (with a plateau at $\tau = 0.935$), potential and kinetic energies is identical to Fig. 1 of Ref. [8]. Thus, the clarification provided in [5] as to the origin of the TPA lacks support within the scalar field cosmology itself. As we shall see below, in the extended scalar field cosmology, that is the 3-fluid cosmology, the energy density need not exhibit a plateau during the TPA as Fig. 1 reveals.

For all solutions discussed below we choose $V_0 = 1$ and $\gamma = 1.5$ and we plot in red (analytic) solutions to the scalar field problem and in blue (numerical) solutions to the 3-fluid problem. Our initial conditions are $U(\tau_0) = U_s(\tau_0)$, $W(\tau_0) = W_s(\tau_0)$ and $W'(\tau_0) = W_s'(\tau_0)$ with $\tau_0 = 10^{-10}$.



Figure 1: Case $\lambda = 2\sqrt{2}$ with $m = 1$. (a) (left): The energy densities $\rho = H^2$ for $C = E = 1$ and $\rho_s = H_s^2$. ρ_s has a plateau around $\tau = 0.65$. ρ exhibits no plateau where the acceleration occurs. (b) (medium): The accelerations $\ddot{a}$ for $C = E = 1$ and $\ddot{a}_s$. $\ddot{a}_s$ is maximum at $\tau = 0.72$. (c) (right): $\ddot{a}$ for $C = E = 100$.

We start by discussing the solution plotted in Fig. 1 of Ref. [8] for which $\lambda = 2\sqrt{2}$. For $C = E = 1$ Fig. 1 (b) shows plots of $\ddot{a}_s$ and $\ddot{a}$ which both exhibit three phases: deceleration-acceleration-deceleration

---

[2] Because of different conventions, the values of (λ_R, V_{0R}) used in [8] are such that $\lambda = \sqrt{6}\lambda_R$ and $V_{0R} = 6V_0$.



Figure 2: Case $\lambda = 3$ with $m = 1$. (a) (left): $\ddot{a}$ for $C = E = 1$ and $\ddot{a}_S$. (b) (medium): $\ddot{a}$ for $C = E = 10$. (c) (right): $\ddot{a}$ for $C = 1, E = 100$.



Figure 3: Case $\lambda = 5$. (a) (left): $\ddot{a}$ for $C = E = 1/100$ and $\ddot{a}_S$. (b) (medium): $\ddot{a}$ for $C = E = 1/10$. (c) (right): $\ddot{a}$ for $C = E = 1$.

(DAD). For larger values of the constants $C = E = 100$ as in Fig. 1 (c), the TPA is right-shifted and the acceleration is smoothed. For much larger C and E the acceleration is completely smoothed, except near the origin, and the solution no longer exhibits the DAD phases; rather, it exhibit an eternal deceleration expansion [an example is shown in Fig. 3 (c) for $\lambda = 5$.] In Fig. 2 we have similar plots for $\lambda = 3$.

These same conclusions extend to the cases $\lambda < 3\sqrt{2}$ where no TPA occurs in both problems. In this case the acceleration curve, which exhibits deceleration-eternal acceleration (DA) expansions, is right and downward shifted with increasing C and E until an eternal deceleration occurs.

Curiously enough, the trigonometric solutions (5) and (6) were not discussed deeply in the literature [8, 14]. Contrary to a statement made in [14] which claims that these solutions exhibit only the DA expansions, we have checked that for $\lambda > 3\sqrt{2}$ up to (and above) $\lambda = 100$ these solutions do indeed exhibit the DAD expansions, as the red plot in Fig. 3 (a) depicts. Moreover, the acceleration period coincides with a plateau for the energy density. Numerical solutions to the 3-fluid problem are plotted in Fig. 3 showing the acceleration for increasing values of the constants. The acceleration phase disappears as the constants increase and the solution ends with an eternal deceleration expansion.

We have checked that the conclusions drawn for the case $\gamma = 1.5$ extend to the case $\gamma = 1.9$ and most probably to all $1 < \gamma \leq 2$.

Discussion. Solutions to the scalar field problem already predict the DAD expansions for the cosmological evolution of the universe for a wide range of λ. Solutions to the 3-fluid problem add new dimensions: The parameters C and E along with λ allow for fine tuning and horizontal and vertical shifting of the acceleration and density curves to fit observational data. Excess in the OM density (C large) or/and the DM density (E large) results in a smooth acceleration curve, except near the origin, exhibiting no DAD phases; rather, the universe would undergo an eternal deceleration expansion.

No solution with two or more TPA's were found. So, the model, with OM and DM included, still fails to predict the early universe inflation.

References

[1] B. Ratra and P.J.E. Peebles, Phys. Rev D. **37**, 3406 (1988).

[2] J.M. Overduin and F.I. Cooperstock, Phys. Rev. D **58**, 043506 (1998), arXiv:astro-ph/9805260v1.

[3] L.P. Chimento, Class. Quantum Grav. **15**, 965 (1998).

[4] P. Townsend and M. Wohlfarth, Phys. Rev. Lett. **91**, 061302 (2003), hep-th/0303097.

[5] R. Emparan and J. Garriga, JHEP05, 028 (2003), hep-th/0304124v2.

[6] A. Kehagias and G. Kofinas, Class. Quantum Grav. **21**, 3871 (2004), gr-qc/0402059v2.

[7] E. V. Linder, Phys. Rev. Lett. **70**, 023511 (2004), arXiv:astro-ph/0402503v1.

[8] J.G. Russo, Phys. Lett. B **600**, 185 (2004), arXiv:hep-th/0403010v3.

[9] E. Elizalde, S. Nojiri and S.D. Odintsov, Phys. Rev D. **70**, 043539 (2004), arXiv:hep-th/0405034v2.

[10] P. Townsend, in XIV-th International Congress on Mathematical Physics, Ed. J.C. Zambrini, World Scientific, Singapore, pp. 655-662 (2005), hep-th/0308149.

[11] A.W. Brookfield et al., Phys. Rev. Lett. **96**, 061301 (2006), arXiv:astro-ph/0503349v2.

[12] J.A.S. Lima, F.E. Silva and R.C. Santos, Class. Quantum Grav. **25**, 205006 (2008), arXiv:0807.3379v2 [astro-ph]

[13] S. Basilakos, M. Plionis and S. Solà, Phys. Rev. D **80**, 3511 (2009), arXiv:0907.4555v2 [astro-ph.CO].

[14] A.A. Andrianov, F. Cannata and A.Yu. Kamenshchik, JCAP10, 004 (2011), arXiv:1105.4515v1 [gr-qc].

[15] M. Tsamparlis and A. Paliathanasis, Class. Quant. Grav. **29**, 015006 (2012), (arXiv:1111.5567v1 [astro-ph.CO].

[16] A.A. Andrianov, F. Cannata and A.Yu. Kamenshchik, Phys. Rev D. **86**, 107303 (2012), arXiv:1206.2828v1 [gr-qc].